

blank page

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



## Pilferer 500

### HERBICIDE

ACTIVE CONSTITUENT: 500g/L PROPYZAMIDE

**GROUP D HERBICIDE**

For selective control of certain grasses and broad leaf weeds in canola, grain legumes, legume pasture, lettuce and turf according to the Directions for use Table.

**IMPORTANT: Read this booklet before use.**

APVMA Approval No. 68267/122248

**AIRR APPARENT PTY LTD**  
15/16 Princes Street, Newport NSW 2106  
M 0411 227 338 | E [enquiries@apparentag.com.au](mailto:enquiries@apparentag.com.au) | [www.apparentag.com.au](http://www.apparentag.com.au)

## DIRECTIONS FOR USE

### Restraints:

Do not apply more than once per season.

Do not apply to water logged soils.

Do not apply to land where runoff to aquatic areas or sensitive crops like cereals is likely.

Crops	Weeds Controlled	State	Rate L/ha
<b>Canola</b>	Annual Ryegrass, Winter Grass, Barley	All States	1.0
<b>Grain Legumes</b> (including lupins, chickpeas, lentils, field peas, faba beans, navy beans)	Grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue, Rats Tail Fescue		
<b>Legume Pastures</b> (including lucerne, clover, medics etc. grown for forage, hay or seed production)		NSW, ACT, Tas, SA, WA only	1.0-1.5
	Wild Oats, Perennial Ryegrass, Bent Grass, English Couch, Yorkshire Fog Grass, Chickweed, Seedling Sorrel, Blackberry Nightshade, Patterson's Curse, Seedling Wireweed		1.5-2.0
Lettuce (direct sown)	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter Grass,	All States	4.5 in 330-550L of water
Lettuce (transplanted)	Rye Grass, Summer Grass, Wireweed, Prince of Wales Feather, Shepherd's Purse, Nettles	Vic only	2.3 in 300-550L of water
Turf and lawns Common Couch	Winter Grass	All States	1.2 (12mL/100m <sup>2</sup> )
Queensland Blue Couch		Qld only	0.6 (6mL/100m <sup>2</sup> )
Buffalo Grass		WA only	1.2 (12mL/100m <sup>2</sup> )

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

Critical Comments	
Apply to moist soil prior to seeding only. Incorporate by sowing (IBS) with knife or blade points. A rainfall event (25mm) between application and sowing, or soon after sowing is required for best results. Avoid seeding practices that move treated soil back into the furrow.	
Heavy rainfall events that wash product back into the furrow can delay emergence and cause crop injury. Do not use on white sandy soils, or paddocks that have been recently mouldboard ploughed or spaded.	
Apply when weeds are at post or pre-emergent stages.	
SEEDLING CROPS: Apply after the third trifoliate leaf is fully developed.	
ESTABLISHED CROPS: Apply any time before the end of July.	
Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended.	
<b>BEST RESULTS: When applied during cool and moist conditions.</b> Rain or irrigation is necessary to thoroughly wet the top 3 to 4cm of soil. This is similar to 25mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting <b>PRIOR</b> to treatment is recommended.	
Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.	
Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.	
Apply on light sandy soils, low in organic matter.	
<b>Rate/Bowling Green</b> 37m x 37m	Apply during the autumn and winter if established turf is infested with seedling winter grass. Ensure a uniform coverage. Irrigate immediately after application with 12-25mm irrigation or rain and again 24 hours later. Repeat treatment if necessary on Qld Blue Couch after 4 weeks. DO NOT RESEED FOR 60 DAYS AFTER TREATMENT. Use <b>ONLY</b> on established turf. <b>DO NOT</b> use on bluegrass, ryegrass, fescue or bent grass lawns.
170mL	
85mL	
170mL	

## **WITHHOLDING PERIODS:**

### **CANOLA:**

**HARVEST – NOT REQUIRED WHEN USED AS DIRECTED.**

**GRAZING – DO NOT GRAZE OR CUT FOR STOCK FOOD 12 WEEKS AFTER APPLICATION.**

### **GRAIN LEGUMES:**

**HARVEST – NOT REQUIRED WHEN USED AS DIRECTED.**

**GRAZING – DO NOT GRAZE OR CUT FOR STOCK FOOD 12 WEEKS AFTER APPLICATION.**

## **ALL OTHER CROPS:**

**DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.**

## **GENERAL INSTRUCTIONS**

Best results are obtained by applying the product to cool, wet soil.

### **MIXING**

Prior to pouring, shake container vigorously, then add the required quantity of Apparent Pflifer 500 Herbicide to water the spray vat while stirring or with agitators in motion. DO NOT mix with other pesticides.

## **WEED RESISTANCE WARNING**

**GROUP D HERBICIDE**

Apparent Pflifer 500 Herbicide is a member of the amide group of herbicides with diverse sites of action. For weed resistance management this is a Group D herbicide. Some naturally occurring weed biotypes resistant to Apparent Pflifer 500 Herbicide and other amide herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Apparent Pflifer 500 Herbicide or other amide herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, AIRR Apparent Pty Ltd accepts no liability for any losses that may result from the failure of Apparent Pflifer 500 Herbicide to control the resistant weeds. Advice as to strategies and alternative treatments that can be used, should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or an AIRR Apparent representative.

## **PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT graze sheep on treated areas or feed to poultry or pigs.

## **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under weather conditions or from spraying equipment that may cause spray drift onto nearby susceptible plants/crops, cropping land or pastures.

### **STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or territory government regulations. DO NOT burn empty containers or product.

**For refillable containers:** Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### **Safety Directions**

Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow-length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

### **First Aid**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

## **SAFETY DATA SHEET**

Additional information is listed in the safety data sheet (SDS). A material safety data sheet for Apparent Pilferer 500 Herbicide is available from AIRR Apparent Pty Ltd on request.

## **CONDITIONS OF SALE**

AIRR APPARENT Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on APPARENT's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of AIRR APPARENT Pty Ltd has any authority to add to or alter these conditions.