



Levistar LV

ORAL DRENCH

ACTIVE CONSTITUENTS:

80 g/L Levamisole Hydrochloride (equivalent to 68 g/L Levamisole)

Claims: For the control of levamisole sensitive Round Worms including Lungworms in Sheep and Cattle.

For the control of levamisole sensitive gastrointestinal roundworms and lungworms in sheep and cattle. Levistar LV contains levamisole, a member of the imidazothiazole (clear) group of anthelmintics (drenches) and is effective in cattle and sheep against: mature and immature Barber's Pole Worm, Stomach Hair Worm, Black Scour Worm, Small Intestinal worm (*Cooperia* spp.), Thin Necked Intestinal Worm, Hookworm, Large Mouthed Bowel Worm, Strongyloides, Nodule and Large Bowel Worms, Large Lungworm and Small Brown Stomach Worm in sheep and mature (adult only) *Ostertagia* spp. in cattle under 12 months of age. Where *Ostertagia* are the problem in cattle over 12 months of age, Starvec Pour-On for Beef and Dairy Cattle is the recommended product.

Controls both susceptible and benzimidazole resistant strains of Barber's pole worm (*Haemonchus* species) and Black scour worm (*Trichostrongylus* species) in sheep.

DOSING AND ADMINISTRATION:

Levistar LV is given as a drench by mouth in cattle and sheep. Dose rates for cattle: 5 mL per 50 kg liveweight.

The live weights should be estimated carefully so that the dose can be calculated accurately to ensure optimum efficiency with economy. The following Cattle and Sheep tables may be used as a guide.

Cattle Dose Rate Table:

LIVEWEIGHT	DOSE VOL (ML)	NO. OF CATTLE TREATED
Up to 50	5	2000
51 - 100	10	1000
101 - 150	15	666
151 - 200	20	500
201 - 250	25	400
251 - 300	30	333
301 - 350	35	285
351 - 400	40	250
401 - 450	45	222
451 - 500	50	200
501 - 550	55	181
551 - 600	60	166
601 - 650	65	153

Dose rates for sheep: 1 mL per 10 kg liveweight.

Always dose on a liveweight basis. Check the accuracy of drenching equipment before and during drenching procedures. The following table may be used as a guide.

Sheep Dose Rate Table:

LIVEWEIGHT	DOSE VOL (ML)	NO. OF SHEEP TREATED
Up to 10	1	10000
11 - 20	2	5000
21 - 30	3	3333
31 - 40	4	2500
41 - 50	5	2000
51 - 60	6	1666
61 - 70	7	1428
71 - 80	8	1250

WITHHOLDING PERIODS

Meat: DO NOT USE less than 3 days before slaughter for human consumption.

Milk: Zero (0) days.

Trade Advice: Export Slaughter Interval (ESI):

DO NOT USE less than 7 days before slaughter for export.

For SDS or technical information go to independentsown.com.au

