

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY

# MOXISTAR®

## LONG ACTING INJECTION FOR SHEEP

**ACTIVE CONSTITUENT: 20 mg/mL MOXIDECTIN**

For the treatment and control of roundworms, nasal bot and itchmite in sheep.  
For protection against severe challenge by *Haemonchus contortus* (Barber's Pole Worm) for up to 4 months.

## 500 mL

Read the carton for full instructions.

### DIRECTIONS FOR USE

#### Restraints:

**DO NOT USE** in ewes, which are producing or may in the future produce milk that may be used or processed for human consumption.

**Re-treatment interval: DO NOT** retreat sheep for at least 91 days after last treatment.

#### Dosage and Administration:

**Use the contents within 6 months of first broaching the vial. Discard the unused portion.**

Dose rate: 1 mg moxidectin/kg liveweight  
(1 mL/20 kg liveweight)

Administer SC at base of the ear or high on the neck behind the ear. Administer the dose according to the dosage table on the carton using an appropriate sheep injector gun. **DO NOT** use in sheep less than 20 kg.

### WITHHOLDING PERIODS

**MEAT: 91 days.**

**MILK: DO NOT USE** in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

### TRADE ADVICE

**EXPORT SLAUGHTER INTERVAL (ESI): 133 days.**

### SAFETY DIRECTIONS

Poisonous if swallowed. May irritate the nose and throat. Will irritate eyes and skin. Avoid contact with eyes and skin. Do not inhale. Wash hands after use.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

### ENVIRONMENTAL PROTECTION

Extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

### STORAGE

Store below 30°C  
(room temperature).  
Protect from light.

BATCH:

EXP:

**May cause respiratory irritation.  
Causes damage to organs through  
prolonged or repeated exposure.**

Do not get in eyes, on skin, or on clothing.

Wash contacted areas thoroughly after handling.

Do not eat, drink or smoke when using this product.

Take off contaminated clothing and wash before reuse.

**IF SWALLOWED:** Rinse mouth. Do NOT induce vomiting.

**IF ON SKIN:** Wash with plenty of soap and water.

Store locked up.

