


# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# Apparent



# Propyzamide 500 WG

## HERBICIDE

ACTIVE CONSTITUENT: 500g/kg PROPYZAMIDE

GROUP **3** HERBICIDE

For selective control of certain grasses and broad leaf weeds in canola, grain legumes, legume pasture, lettuce and turf according to the Directions for use Table.  
**IMPORTANT: Read the attached leaflet before use.**

PROUDLY



OWNED



The Choice is  
**Apparent**



NET CONTENTS

15

KILOGRAMS

AIRR APPARENT PTY LTD

15/16 Princes Street, Newport NSW 2106

P 03 5820 8400 | E enquiries@apparentag.com.au | www.apparentag.com.au

# Apparent Propyzamide 500 WG HERBICIDE

### DIRECTIONS FOR USE

#### Restraints

Do not apply more than once per season.  
Do not apply to water logged soils.  
Do not apply to land where runoff to aquatic areas or sensitive crops like cereals is likely.

Crops	Weeds Controlled	State	Rate/ha	Critical Comments
<b>Canola</b> <b>Grain Legumes</b> (including lupins, chickpeas, lentils, field peas, faba beans, navy beans)	Annual Ryegrass, Winter Grass, Barley Grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue, Rats Tail Fescue	All States	1.0kg	Apply to moist soil prior to seeding only. Incorporate by sowing (IBS) with knife or blade points. A rainfall event (25mm) between application and sowing, or soon after sowing is required for best results. Avoid seeding practices that moves treated soil back into the furrow. Heavy rainfall events that wash product back into the furrow can delay emergence and cause crop injury. Do not use on white sandy soils, or paddocks that have been recently mouldboard ploughed or spaded.
<b>Legume Pastures</b> (including lucerne, clover, medics etc. grown for forage, hay or seed production)	Wild Oats, Perennial Ryegrass, Bent Grass, English Couch, Yorkshire Fog Grass, Chickweed, Seedling Sorrel, Blackberry Nightshade, Patterson's Curse, Seedling Wireweed	NSW, ACT, Tas, SA, WA only	1.0-1.5kg	Apply when weeds are at post or pre-emergent stages. SEEDLING CROPS: Apply after the third trifoliate leaf is fully developed. ESTABLISHED CROPS: Apply any time before the end of July.
Lettuce (direct sown) Lettuce (transplanted)	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter Grass, Rye Grass, Summer Grass, Wireweed, Prince of Wales Feather, Shepherd's Purse, Nettles	All States	4.5kg in 330-550L of water	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.
Turf and lawns Common couch	Winter Grass	All States	2.3kg in 300-550L of water	Apply on light sandy soils, low in organic matter.
Queensland blue couch		Old only	1.2kg (12g/100m <sup>2</sup> )	<b>Rate/Boiling Green</b> 37m x 37m 170g 85g
Buffalo grass		WA only	0.6kg (6g/100m <sup>2</sup> ) 1.2kg (12g/100m <sup>2</sup> )	170g

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**  
**WITHHOLDING PERIODS:**

#### CANOLA:

**HARVEST – NOT REQUIRED WHEN USED AS DIRECTED.**

**GRAZING – DO NOT GRAZE OR CUT FOR STOCK FOOD 12 WEEKS AFTER APPLICATION.**

#### GRAIN LEGUMES:

**HARVEST – NOT REQUIRED WHEN USED AS DIRECTED.**

**GRAZING – DO NOT GRAZE OR CUT FOR STOCK FOOD 12 WEEKS AFTER APPLICATION.**

#### ALL OTHER CROPS:

**DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.**

#### GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet soil.

#### RESISTANCE WEED WARNING

##### GROUP 3 HERBICIDE

Apparent Propyzamide 500 WG Herbicide is a member of the amide group of herbicides with diverse sites of action. For weed resistance management this is a Group 3 herbicide. Some naturally occurring weed biotypes resistant to Apparent Propyzamide 500 WG Herbicide and other amide herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Apparent Propyzamide 500 WG Herbicide or other amide herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, AIRR APPARENT Pty Ltd accepts no liability for any losses that may result from the failure of Apparent Propyzamide 500 WG Herbicide to control the resistant weeds.

Advice as to strategies and alternative treatments that can be used, should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or AIRR APPARENT representative.

#### Spray Preparation

Apparent Propyzamide 500 WG Herbicide is a Wettable Granule formulation to be blended with water and applied as a spray. Granule formulations are designed to be added to large volumes of water and dispersed slowly under agitation. When using larger boomsprays and unable to add granules directly to a large volume in the tank via the top of the boomspray, pre-mixing may be necessary as follows:

- Partly (30%) fill the boomspray tank with water.
- Slowly add required amount of Apparent Propyzamide 500 WG Herbicide for the area to be sprayed into Granni Pot® or other pre-mixing vessel while agitating well.
- Ensure high water volumes and DO NOT add to the spray tank until the pre-mix is fully dissolved. This may take several mixes to ensure the correct amount is added to the boomspray tank.
- Add the pre-mixed Apparent Propyzamide 500 WG Herbicide into the boomspray tank with the agitation system operating.
- Fill up to the correct volume with water.
- The mixture must be kept in suspension at all times by continuous agitation.
- Tank-mixtures should be tested for physical compatibility and efficacy before use.

DO NOT mix with MCPA. If tank mixing, Apparent Propyzamide 500 WG Herbicide must be in suspension before adding other herbicides, insecticides or surfactants. Wash sprayer thoroughly after use.

#### Equipment

Always ensure the sprayer is clean, using a suitable spray and boom cleaner as directed. It is recommended that for turf situations, up to 100L/ha spray volume is required.

#### COMPATIBILITY

Apparent Propyzamide 500 WG Herbicide is compatible with a range of knockdown and pre-emergent herbicides and insecticides. The following table provides some recommendations on the suitability of tank mix combinations.

Physical compatibility with Apparent Propyzamide 500 WG Herbicide should be determined prior to mixing with a product not listed in this table, or when mixing Apparent Propyzamide 500 WG Herbicide as a component of a 3-way tank mix.

Always read the product label for the manufacturer's tank mix recommendations to determine individual product compatibility options and correct mixing orders for individual products. If unsure, perform a jar test before proceeding to determine physical compatibility. Physical compatibility does not always guarantee biological compatibility.

Herbicides	Compatibility recommendation
Apparent Atrazine 900 WG	Compatible
Apparent Diuron 900 WG	Compatible
Ecopar (pyratufen-ethyl)	Compatible
Glyphosate present as the potassium salt based products**	Not recommended - incompatible
Apparent Glyphosate 450	Conditional
Glyphosate Dry WG680	Compatible
Apparent Haloxypop 520	Compatible
Apparent Clobberlime 400	Conditional
MCPA products	Not recommended - incompatible
Apparent Oxyfluorelone 240 EC	Conditional
Apparent Paraquat 250 SL	Not recommended - incompatible
Paraquat 360 SL	Not recommended - incompatible
Apparent Disturb 875 WG	Compatible
Apparent Trifluralin 480 EC	Not recommended - incompatible
Insecticides	Compatibility recommendation
Dimethoate 400 EC	Not recommended - incompatible
Dimethoate 290 SL	Compatible
Apparent Annihilate 100	Compatible
Apparent Stockade 250 EC	Conditional

#### Explanatory notes on recommendations

**Compatible:** Products considered physically compatible with Apparent Propyzamide 500 WG in a two-way mix.

**Conditional:** Tank mixes of Apparent Propyzamide 500 WG with the following product may result in nozzle or filter blockage if vigorous agitation in the spray mix unit is not maintained during the entire spray operation.

**Not recommended - incompatible:** Tank mixes of Apparent Propyzamide 500 WG with the following product is likely to form a slimy residue which can block spray nozzles and filters during application.

\*\* It is not recommended to mix Apparent Propyzamide 500 WG with any products containing **Glyphosate present as the potassium (K) salt**.

Information on compatibility is understood to be correct at the time of publication, however products may vary from time to time.

Additional information required under the Globally Harmonised System (GHS) classification of the substance/mixture:

**Suspected of causing cancer. Very toxic to aquatic life. May cause long lasting harmful effects to aquatic life.**

Avoid release to the environment. IF ON SKIN: Wash with plenty of soap and water. IF exposed or concerned: Get medical advice/attention: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If skin irritation occurs: Get medical advice/attention. If eye irritation persists: Get medical advice/attention. Take off contaminated clothing and Wash before reuse. Collect spillage. Store locked up.

APVMA Approval No: 80689/122250

Batch No:

Date of Manufacture:

200



Scan QR code to view booklet on mobile device

IN A TRANSPORT EMERGENCY  
**DIAL 000**  
POLICE OR FIRE BRIGADE



9 345724 008446